

<b>REPORT DOCUMENTATION PAGE</b>				Form Approved OMB No. 0704-0188	
<small>The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.</small> <b>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</b>					
1. REPORT DATE (DD-MM-YYYY) 17/Jan/2002		2. REPORT TYPE MAJOR REPORT		3. DATES COVERED (From - To)	
4. TITLE AND SUBTITLE REVASCULARIZATION FOR CHRONIC MESENTRIC ISHEMIA COMPARISON OF ENDOVASCULAR AND OPEN SURGICAL INTERVENTION				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) MAJ BOWSER ANDREW N				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) UNIVERSITY OF SOUTH FLORIDA				8. PERFORMING ORGANIZATION REPORT NUMBER CI02-4	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) THE DEPARTMENT OF THE AIR FORCE AFIT/CIA, BLDG 125 2950 P STREET WPAFB OH 45433				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Unlimited distribution In Accordance With AFI 35-205/AFIT Sup 1				DISTRIBUTION STATEMENT A: Approved for Public Release - Distribution Unlimited	
13. SUPPLEMENTARY NOTES					
<div style="font-size: 2em; font-weight: bold;">20020204 103</div>					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES 1	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			19b. TELEPHONE NUMBER (Include area code)

## ABSTRACT:

**Purpose:** To compare the patient groups and outcome following treatment of symptomatic chronic mesenteric ischemia using percutaneous angioplasty/stenting (PAS) or open surgical reconstruction (OS).

**Methods:** Over a 9-year period, 40 patients (23 women, 17 men) presenting with atherosclerotic occlusive (ASO) disease involving the visceral arteries underwent PAS (18 patients; 20 vessels treated) or OS (22 patients; 35 vessels treated). Patient demographics, extent of visceral artery revascularization, and procedure outcome were compared ( $\chi^2$ , t-test, or life-table analysis) between the two treatment groups.

**Results:** The PAS and OS patient groups had similar ( $p>0.1$ ) duration of symptoms, weight loss, pre-operative serum albumin, and ASO risk factors. Patients selected for PAS were older (68 versus 59.5 years,  $p=0.03$ ) and had more advanced active cardiac disease. Endovascular intervention involved balloon angioplasty and stenting of only one artery in all but two patients (SMA 9/18, 50%; celiac 6/18, 33%; IMA 1/18, 6%; SMA & celiac 2/18, 11%), while the majority (13/22, 59%) of OS patients had 2-vessel revascularization by bypass alone ( $n=8$ ), bypass & endarterectomy ( $n=3$ ), or endarterectomy alone ( $n=2$ ). 30-day procedural failure (death due to multisystem organ failure, vessel re-intervention) was similar in the PAS (mortality 12% [2/18], PTA site stenosis, 1/18) and OS (mortality 9% [2/22], vessel occlusion 0/22) groups. Mean duration of ICU care/hospitalization was higher ( $p<0.01$ ) for OS (8/24 days) patients than PAS (1.5/6 days) patients and related to intestinal resection (4 patients) and slower return of GI function, but overall morbidity (wound/PTA site, cardiac, pulmonary, renal) was comparable (PAS, 44%; OS, 45%). Overall survival at 3 years was similar (OS, 88%; PAS, 76%) and no patient required late gut resection for ischemia. Primary patency @ 14 months (No. of vessels with  $<50\%$  stenosis/No. of vessels) assessed by visceral duplex/angiography was 51% for PTA- versus 82% for surgical-treated patients. During follow-up, more surviving PAS patients (5 of 16) required intervention for recurrent symptoms or asymptomatic stenosis compared to OS (2 of 20) patients. Six of 7 re-interventions involved PTA of a stent or an additional visceral artery. With re-intervention, all patients became symptom free.

**Conclusions:** Treatment of chronic mesenteric ischemia by endovascular or standard surgical revascularization produced similar outcomes, despite differences in patient age, severity of gut ischemia, and need for re-intervention. Endovascular treatment should be limited to high-risk patients in whom single-vessel PTA restores gut blood flow. Surveillance after intervention is recommended since one-quarter (7 of 36) of patients developed recurrent symptoms or asymptomatic re-stenosis – the majority of which were amenable to endovascular treatment.